

Tropical ecology

WBNZ-849

starting 14:45 (as in USOS)

Ryszard Laskowski

Institute of Environmental Sciences, Jagiellonian University

<http://www.cyfronet.krakow.pl/~uxlaskow/>

1. About the course
2. Lecture #1: Introduction to tropical ecology

Course organization

- **Place:** Institute of Environmental Sci., Room 1.1.1
- **Time:** Friday, 14:45 – 17:15
 - 8 x 3 h (lectures & discussion classes)
 - 2 seminars (3 h each)
- **Teachers:** Marcin Czarnołęski, Wojciech Fiałkowski, Paweł Koteja, Ryszard Laskowski, Krzysztof Wiąckowski
- **Evaluation:**
 - final exam (5-6 open questions): 80%
 - active participation in classes: 20%




Teachers' emails

- marcin.czarnoleski@uj.edu.pl
- wojciech.fialkowski@uj.edu.pl
- pawel.koteja@uj.edu.pl
- ryszard.laskowski@uj.edu.pl
- krzysztof.wiackowski@uj.edu.pl



Reading

- Articles and textbooks available at the course website

- Books from the Library of Natural Sciences



Supplementary reading in Polish



ATTENTION:

The 'Tropical Ecology' course (WBNZ 849) is the prerequisite for 'Tropical Ecology Field Course' (WBNZ 850)

Topics:

Introduction to tropical ecology: tropical biomes – geographical distribution and characteristics

Destruction and protection of tropical ecosystems

Equatorial rainforests – the most diverse biome on Earth

- gradients in biodiversity and theories explaining them
- diversity in life strategies

Adaptations in animals to hot deserts

Biology of coral reefs and mangroves: environmental conditions and biodiversity.

Introduction to tropical ecology


Where are the tropics?

- **Origin of the term:** from Greek *τρόπος* (*tropos*) = *turn* (the sun appears to "turn back" at the solstices)
- ➔ Area between the *Tropic of Cancer* ($23^{\circ}30'N$) and the *Tropic of Capricorn* ($23^{\circ}30'S$)
- ➔ Area of the Earth where the Sun is 90° above the horizon at least once every year
- ➔ = *tropical zone* = *torrid zone*


Where are the tropics located?

The simplest possible answer:

The area between the Tropic of Cancer and the Tropic of Capricorn



Why should we study tropical ecology?




Species richness on Earth (per $1^\circ \times 1^\circ \approx 9274 \text{ km}^2$)

Grenyer, R. et al. 2006. Global distribution and conservation of rare and threatened vertebrates. *Nature* 444: 93-96. (pdf available for course participants at the course web page)

Species richness in tropics

Taxonomic group	Poland (312 000 km²)	Uganda (241 000 km²)
vascular plants	2700	4900
mammals	109	330
birds	446	1061
reptiles	9	165
amphibians	18	52


Biodiversity hotspots on Earth



„As many as 44% of all species of vascular plants and 35% of all species in four vertebrate groups are confined to 25 hotspots comprising only 1.4% of the land surface of the Earth.”

Myers, N. et al. 2000. Biodiversity hotspots for conservation priorities. Nature 403:853-858. (pdf available for course participants at the course web page)

Tropical climates according to Wladimir Köppen



Tropical climates according to Köppen


■ Group A: Tropical (megathermal) climates

- *Af*: Tropical rainforest climate
(~ 5 - 10° of the equator; in coastal areas can extend to 25°; no seasonality) =
hygromegathermal
- *Am*: Tropical monsoon climate (further from the equator; two seasons – rain and dry)
- *Aw*: Tropical savanna climate (two seasons, wet and dry – very clear and pronounced)

■ Group B: Dry climates (arid and semiarid)

- Only partly belong to tropics


1. Tropical rainforests




Climatic diagram for Belem (Brazil)

Tropical rainforests: geographical distribution


around equator (ca. 10°S – 10°N)



Iquitos, Peru



Manaus, Brazil



Tropical rainforests: characteristics

- Very high annual rainfall: **at least 1700 – 2000 mm**
- Average annual temperature: **27 – 30°C**
- High rate of biogeochemical cycles
- Soils: low in organic matter and nutrients due to intensive weathering (**laterization → oxisols**)
- **Four-layer forests**: (1) emergent layer – single trees above the canopy (60-70 m); (2) canopy layer (30-45 m); (3) understory layer (only ca. 5% of light!); (4) forest floor (only ca. 2% of light)
- Richness of epiphytes and lianas
- Extreme species richness: **>30% of all plant and animal species living on Earth** at only **6%** of Earth surface!

Tropical rainforests: types

- **Lowland equatorial evergreen rainforests**
 - annual precipitation above 2000 mm
 - Amazon, Orinoco and Congo basins, Indonesia, New Guinea
- **Wet broadleaf forests partly evergreen**
 - high annual rainfall, warm and wet summer and cooler and dryer winter
 - Central America, Caribbean, West Africa, India, Indochina
- **Montane cloud forests**
 - cooler mountain climate, high rainfall, low cloud cover
 - tropical and subtropical mountains
- **Floodplain forests**
 - environmental conditions similar to lowland evergreen forests but in poorly drained areas → flooding
 - Borneo, Sumatra, Malay Peninsula, Indochina

Nutrient turnover rate

Average retention time of dead organic matter and nutrients in forest litter:
boreal forest (taiga), temperate broadleaf forest, and equatorial rainforest
(*time in years*)

Biome	Organic matter	N	P	K	Ca	Mg
Taiga	353	230	324	94	149	455
Temperate forest	4	5.5	5.8	1.3	3.0	3.4
<i>Rainforest</i>	<i>0.4</i>	<i>2</i>	<i>1.6</i>	<i>0.7</i>	<i>1.5</i>	<i>1.1</i>

Productivity and carbon accumulation


Average NPP of selected biomes ($\text{kg x m}^{-2} \text{ x year}^{-1}$),
carbon accumulation rate ($\text{g x m}^{-2} \text{ x year}^{-1}$) and C(biomass)/C(soil)

Biome	Productivity	C accumulation rate	C(b)/C(s)
Taiga	0.8	11.7 – 15.3	0.55
Temperate forest	1.2	0.7 – 5.1	1.13
<i>Rainforest</i>	2.2	2.3 – 2.5	1.68

Main carbon pools in primeval tropical rainforests

Part of the ecosystem	Accumulated carbon (t C/ha)
Alive plants (above and underground)	210
Dead trees and litter	10
Soil	100
<i>TOTAL:</i>	<i>320</i>


Tropical rainforest soils



Ferrasols (FAO) = Oxisols (USDA): location and pedogenesis

- Earlier called *laterites*; acc. to FAO - *ferrasols*
- Definition: soils containing in the whole profile $\leq 10\%$ leachable materials and $< 10\%$ base saturation; high content of Fe and Al oxides
- Location: ca. 1/3 of the Earth's continental land area, mostly 15-25°S – 15-25° N
- Pedogenesis – tropical weathering (*laterization*):
 - high precipitation + $\text{CO}_2 \rightarrow$ chemical weathering and leaching of humic materials and minerals from the soil profile
 - only stable Fe i Al oxides remain \rightarrow rusty-red color

Laterization



Ferrasols (Oxisols) – Kenya



Laterization – consequences:

- Leaching of virtually all organic matter and nutrients
 - soils very poor in nutrients
 - very small reservoirs of soil organic matter
 - plants have to use (re-cycle) all minerals released from decomposing litter very efficiently
 - no nutrient supply after forest destruction and removal of plants → soils become infertile very quickly → difficult forest regeneration
 - primeval forests replaced with secondary ecosystems (secondary forests of bushes)

Four-layer forest structure



Emergent layer:

60-70 m high broadleaf trees

birds, monkeys, bats, butterflies



Canopy layer

30-45 m high, very dense layer

toucans, snakes, treefrogs, beetles



Understory layer

3-4 m high, little sunshine

treefrogs, beetles, leopards/jaguars



Forest floor

very dark → few plants

anteaters, snakes, frogs, beetles

Four-layer forest structure



Diversity of life forms: epiphytes and vines



Extreme species richness



Species richness of tropical rainforests

- At 10 ha of forest in Borneo – up to 700 tree species → as many as in whole N. America!
- At 1 Peruvian tree – 43 ant species → as many as in whole UK!
- Ca. 3000 fish species in the Amazon river – more than in whole North Atlantic ocean!
- Species numbers at 15 km² in Costa Rica:
 - mammals – 117 (*in whole Poland 105*); birds – 410 (*435*); reptiles – 86 (*9*); amphibians – 43 (*18*); moths – 4000 (*1200*); vascular plants – 1668 (*2700*)

Tropical rainforests: montane cloud forests (fog forest)

- Specific type of tropical rainforests:
 - area: tropical mountains
 - environmental conditions: persistent or frequent low-level cloud cover and fog → reduction of direct radiation and evapotranspiration, very high humidity
 - ecosystem characteristics: particularly rich in epiphytes (mosses, ferns, orchids, etc.)

Montane cloud forests




Africa



Venezuela




Ecuador



Ecuador



2. Tropical and subtropical seasonal dry broadleaf forests (monsoon forests)



Climatic diagram for Calcutta (India)

Monsoon forests: geographical distribution

Two belts N and S from equatorial rainforests: ca. 10° – 20°N & 10° – 20°S



Monsoon forests: characteristics

- High average annual temperature
- High annual rainfall (~1000 – 2000 mm/year)
- Clearly pronounced, long (few months) dry season
 - ➔ most trees shed leaves in dry season;
 - ➔ plants accumulating water;
 - ➔ rich understory layer (plenty of sunlight in dry season)
 - ➔ three layers: (1) tree canopy; (2) understory; (3) forest floor

Main carbon pools in monsoon forests

Ecosystem part	Accumulated carbon (t C/ha)
Alive plants (above- and underground)	150
Dead trees and litter	10
Soil	100
<i>TOTAL:</i>	<i>260</i>

Monsoon forest in dry season



Monsoon forest in Trinidad

Monsoon forests: characteristic tree species




Teak tree
(*Tectona sp.*)



Ebony tree (*Diospyros sp.*) 39/62

Tropical grasslands



Climatic diagram for Nairobi (Kenya)


Tropical grasslands in the world

- Africa:
 - **Savannah**, e.g. Serengeti, Masai Mara – high grasses with scattered acacia trees; large herbivores (40 ungulate species) and carnivores
- South America:
 - **Llanos** in Venezuela (Orinoco basin) – flooded every year, with gallery forests
 - **Cerrado** in Brazil – grassland covered with forest of different density and gallery forests; high plant diversity
- Australia:
 - **Savannah** (Northern Australia) – grassland with scattered eucalyptus trees; herbivores – kangaroos and man-introduced ungulates

Savannah

- Average annual precipitation 1000-1500 mm (Köppen's *Aw* climate)
- Distinct, long dry season;
- Temperature: 20-30°C
- NPP: ca. 0.7 kg d.w. m⁻² year⁻¹
- Plants – adaptations
 - to dry season: deep tap roots, thick bark, shedding leaves, storage organs (mostly underground)
 - to herbivores: solid sharp leaves, bitter taste, growing from beneath)

Savannah soils



Tropical grasslands



Savannah (Kenya)



Llanos (Venezuela)



Cerrado (Brazil)



Savannah (Australia)


Main carbon pools in tropical savannah

Ecosystem part	Accumulated carbon (t C/ha)
Alive plants (above- and underground)	35
Dead trees and litter	0
Soil	55
<i>TOTAL:</i>	<i>90</i>

Main carbon pools in tropical grasslands besides savannah

Ecosystem part	Accumulated carbon (t C/ha)
Alive plants (above- and underground)	12
Dead trees and litter	0
Soil	42
<i>TOTAL:</i>	<i>54</i>


Tropical deserts



Climatic diagram for Cairo (Egypt)

Desert soils – aridisols (USDA)

(FAO: gypsisols, calcisols, solonchaks, solonetztes)




Desert soil profile: clearly seen calcareous layer

- Main process: CaCO_3 and MgCO_3 accumulation → development of calcareous layer
 - rain + atmospheric CO_2 → weak carbonic acid
 - dissolving Ca and Mg salts from surface minerals
 - transport to deeper soil layers
 - evaporation → increasing concentration of dissolved minerals
 - solidification of salts from the solution
 - **concentrations of salts toxic to plants and animals**
 - **water-impermeable carbonate layer**


Main carbon pools in tropical deserts

Ecosystem part	Accumulated carbon (t C/ha)
Alive plants (above- and underground)	1
Dead plants and litter	0
Soil	0
<i>TOTAL:</i>	<i>1</i>

Mountains change everything: zonation and characteristics of the vegetation of Mt. Kenya




Mountains change everything: Altitudinal vs latitudinal zonation




Aiba, S. 2016. Vegetation zonation and conifer dominance along latitudinal and altitudinal gradients in humid regions of the Western Pacific (In: Structure and Function of Mountain Ecosystems in Japan, ed. Gaku Kudo).

Mountains change everything: Altitudinal vs latitudinal zonation



Species richness decreases with altitude



Relationship between the number of species and altitude:
bats in Manu Biosphere Reserves (Peru)

Ecofloristic zones in tropical mountains

- **Alpine:** ~3800 – ~4500 m
 - high mountain steppe: Afro-alpine, paramo, puna
- **Subalpine:** ~3400 – 3800 m
 - few lianas and vascular epiphytes, rich moss and lichen flora; characteristic groups: Ericaceae, Brunelliaceae, Asteraceae...
 - 'elf forests' at ridges
- **Montane:** ~2400 – 3400 m
 - short trees, even fewer species; few lianas, still many epiphytes; can be seasonal
- **Submontane:** ~1000 – 2400 m
 - forest similar to that at lower elevation but with fewer species; trees ca. 25-30 m

Tropical mountain zones in the Andes: Venezuela, Pico Bolivar



ca. 4000 m

„Tierra fria”
Paramo



ca. 3000 m

„Tierra fria”
Upper montane
forest



ca. 2000 m


„Tierra templada”
Lower montane forest

Tropical mountain zones in Africa: Tanzania, Kilimanjaro



ca. 4000 m

Alpine



ca. 3000 m

Montane



ca. 2000 m

Submontane

Other tropical plant communities

■ Mangroves

- areas: shallow, muddy sea coasts;
- structure: trees or shrubs, very few or even just one species; no understory and forest floor; few epiphytes and lianas

■ Gallery forests

- areas: along valleys with surface or underground streams
- structure: trees or bushes of different density; possible lianas, few epiphytes

Mangroves



Caribbean coast,
Venezuela



Gallery forests



Gran Sabana, Venezuela

Gallery forests



Gran Sabana,
Venezuela



Example topics for the seminar:

- 1) The newest data on the role of tropical rainforests in global carbon balance.
- 2) The highest tree species diversity in the world – where and why?
- 3) Species diversity (of selected groups) on altitudinal gradient in the tropics.
- 4) Is it possible to restore destroyed tropical rainforests? Área de Conservación Guanacaste – a case study in Costa Rica.
- 5) Tropical diseases: most important diseases, prevention & problems.

...?

Important dates (on my website):

1. 18.10.2024 R. Laskowski – lecture/discussion class: Course plan and rules; Introduction to tropical ecology: tropical biomes – area, climate, soils and characteristics; latitudinal and altitudinal zonation
2. 25.10.2024 R. Laskowski – lecture/discussion class: Anthropogenic destruction and protection of tropical ecosystems; REDD initiative
3. 08.11.2024 K. Wiąckowski – lecture/discussion class: Tropical biodiversity: Latitudinal diversity gradient
4. 15.11.2024 K. Wiąckowski – How can so many species coexist in a tropical rainforest?
5. 22.11.2024 W. Fiałkowski – lecture/discussion class: Biology of coral reefs and mangroves: environmental conditions, biodiversity
6. 29.11.2024 P. Koteja – lecture/discussion class: Adaptations to hot deserts: water balance, behavioural and physiological mechanisms for water conservation; behavioural and physiological thermoregulation, life histories
7. 06.12.2024 M. Czarnołęski – lecture/discussion class: Biodiversity in tropics: diversity in life strategies
8. 13.12.2024 M. Czarnołęski – lecture/discussion class: Tropical societies
9. 10.01.2025 All teachers – seminar (groups 1, 2, 3); Due to the large number of students, there will be parallel seminar sessions.
10. 17.01.2025 All teachers – seminar (groups 4, 5, 6)

Seminar topics to RL: deadline 16th December